Keio University Global Research Institute



KGRI KGRI Lecture Series

Structural and functional abnormalities in cancer and psychiatric diseases



Speaker: Professor Per Uhlén

Dept of Medical Biochemistry and Biophysics Karolinska Institute, Stockholm, Sweden Coordinator: Professor Masato Yasui コーディネーター: 安井正人 医学部薬理学教授 2017年10月13日(金)17:30-18:30 Friday, October 13, 2017 信濃町キャンパス総合医科学研究棟6階 **Center for Integrated Medical Research 6F** 会議室6 Meeting Room 6

Abstract:

Intratumoral heterogeneity is a critical factor when diagnosing and treating patients with cancer. Marked differences in the genetic and epigenetic backgrounds of cancer cells have been revealed by advances in genome sequencing, yet little is known about the phenotypic landscape and the spatial distribution of intratumoral heterogeneity within solid tumors. Our research group has developed a new diagnostic pipeline termed DIPCO, using three-dimensional light-sheet microscopy of cleared solid tumors, to identify structural abnormalities of phenotypic heterogeneity, in the epithelial-to-mesenchymal transition and in angiogenesis, at single-cell resolution in whole formalinfixed paraffin-embedded biopsy samples. We also show that the DIPCO pipeline can identify structural abnormalities to determine tumor stage and stratify patient prognosis from clinical samples with higher accuracy than current diagnostic methods.

The CACNA1C gene is a well-established candidate risk gene in multiple psychiatric disorders, including bipolar disorder, schizophrenia and major depression, and it encodes a voltage-gated Ca2+ channel. However, the mechanism by which the CACNA1C gene imposes its risk is not known. Our research group has used transgenic mice to show that deletion of Cacna1c results in signs of increased anxiety. Magnetic Resonance Imaging scans of these animal revealed structural abnormalities in the brain anatomy. Examining their spontaneous Ca2+ activity showed functional abnormalities that was confirmed by mathematical modeling. We hypothesize that spontaneous Ca2+ activity is driven by functional pacemaker cells, expressing slightly more voltage-gated Ca2+ channels than non-pacemakers.

No prior registration is necessary. 事前登録不要

