

機械工学セミナー Mechanical Engineering Seminar

第10回 No. 10

共催：慶應義塾大学理工学部機械工学科
慶應義塾大学グローバルリサーチインスティテュート(KGRI)
Department of Mechanical Engineering, Keio University
Keio University Global Research Institute

日時(Date):

January 11, 2018 (Thu.) 16:45~18:00

場所(Venue):

14棟創想館2階 セミナールーム3 (14th Building 2F Seminar room 3)

講演題目(Title):

Engineering Approaches of Medical Images

講演者(Speaker):

Taeyong Lee, Associate Professor
Division of Mechanical and Biomedical Engineering
Ewha Womans University, Republic of Korea



Abstract:

Osteoporosis is a prevalent problem amongst the elderly. Bone mineral density (BMD) obtained from dual X-ray absorptiometry (DXA) is the golden standard in diagnosing osteopenia ($-1.0 < t < -2.5$) and osteoporosis ($t > -2.5$). However, following osteoporosis therapy, increases in BMD may be unreliable. Although hip fracture risk can be reduced with the aid of drugs, treated patients still face considerable risk as most people who sustain hip fracture do not have generalized osteoporosis. A study of the local distribution of bone mass was necessary as they contribute to the geometry and consequently the bone strength. By identifying the respective regions in the femoral neck, the geometric changes were localized and differed between each patient, proving that drug treatment elicits local changes in Rmean and CTmean. Numerical analysis also validated the above findings, where critical strain regions were predicted at similar zones and this is coherent with the fact that reduced thickness of the cortical bone has been related to increased risk of fracture initiation. Hence, from individual radar plots, we can determine if the effect of drugs had outweighed the effect of aging. We can then propose a course of treatment drug better suited for the patient in the clinical scenario. Clinically, little conclusion can be drawn from just the BMD in osteopenic/osteoporotic patients. This emphasizes the necessity of using geometry and structure to predict fracture risk. Focusing on a patient specific analysis at a local level will improve diagnosis of osteoporosis and ultimately fracture prediction.



お問合せcontact: 植田利久 Toshihisa Ueda , Email: ueda@mech.keio.ac.jp phone: 045-566-1496